

No time machines in classical general relativity.

S Krasnikov

Central Astronomical Observatory at Pulkovo, St.Petersburg, 196140, Russia

E-mail: redish@pulkovo.spb.su

Abstract. Irrespective of local conditions imposed on the metric, any extendible spacetime U has a maximal extension containing no closed causal curves outside the chronological past of U . We prove this fact and interpret it as impossibility (in classical general relativity) of the time machines, insofar as the latter are defined to be causality-violating regions created by human beings (as opposed to those appearing spontaneously).

PACS numbers: 04.20.Gz

1. Introduction

In this paper we prove a theorem which, in physical terms, says that *in classical general relativity a time machine cannot be built*. To formulate the theorem and to substantiate such its interpretation we need some preliminary discussion.

Suppose one wants to undertake a time trip. A possible strategy would be just to look for a ready-made closed timelike curve (CTC), or to wait passively until such a curve appears.

1. Remark. It should be stressed that such expectation is not hopeless however innocent the spacetime looks at the moment. The possibility of a ‘sudden’, ‘unprovoked’ appearance of a CTC, in my view, must be taken quite seriously, neither theoretical, nor observational evidence against them being known. Consider, for example, the Deutsch-Politzer (DP) space [1], which is the spacetime obtained from the Minkowski plane by making cuts along the segments $\{t = \pm 1, -1 \leq x \leq 1\}$ and gluing then the upper bank of each cut to the lower bank of the other cut (see figure 1a). An observer located, say,




Figure 1. (a) The DP space. Curves 1 and 2 are actually continuous. Through each point of the shadowed region pass closed and self-intersecting causal curves (such as 1). (b) The extension C of the Misner space. Causality is violated in the shadowed region.

at a point $x = 0, t = -5$ and fully informed about the geometry of the world at $t < -5$ cannot foretell whether the spacetime will evolve in the Minkowski plane (preserving thus causality) or in the DP space, both, in particular, being (in the four-dimensional case) the solutions of the Einstein equations with the same (zero) source.

Still, the *discovery* of a ready-made CTC is a matter of luck. The alternative would be the *creation* of such a curve. In particular, manipulating with matter — and thus according to the Einstein equations with the metric — an advanced civilization could try to *force* the spacetime to evolve into a time machine. It is the conjecture that, given a suitable opportunity, the civilization can succeed [2], that initiated intensive studies of

the time machine (see [3] for reviews and references). The difference between ‘finding’ a closed causal curve (or ‘causal loop’ for brevity) ℓ and ‘manufacturing’ it is the central point of our consideration. In distinguishing these two possibilities, I proceed from the idea[‡] that for the existence of ℓ to be attributable to the activity that took place in a region U , ℓ and U must satisfy at least the following two conditions:

- (i) ℓ lies in the future of U , and not in its past;
- (ii) a causal loop satisfying (i) exists in *any allowed maximal* extension of U .

Correspondingly, if no such U can be found I consider ℓ as ‘spontaneous’ rather than ‘manufactured’ by anybody.

The first of the conditions is self-evident, but the emphasized words in (ii) need some comment. An extendible spacetime U typically has infinitely many maximal extensions (e. g. if U is the Minkowski half-plane $t < 0$ those are the Minkowski plane, the DP space, any plane with the metric $ds^2 = (1 + \omega^2)(dx^2 - dt^2)$, where $\omega(t < 0) = 0$, etc.). However, they are not equipollent from the point of view of general relativity, since in that theory often only those spacetimes are considered as possible models of the Universe which satisfy some local conditions.

2. Definition. We call a condition C *local* if the following is true: C holds in a spacetime M if and only if it holds in any U which is isometric to an open subset of M .

Ideally — that is, if we could describe all existing matter by a single Lagrangian — the local condition required by relativity would be the Einstein equations with the stress-energy tensor in the right hand side corresponding to that Lagrangian. Of course we do not know the Lagrangian and so in practice different more simple local conditions (based, in particular, on our guesses about its properties) are imposed. Examples are

$$G_{\alpha\beta} = -\Lambda 1_{\alpha\beta}, \quad \text{or} \quad G_{\alpha\beta} t^\alpha t^\beta \geq 0 \quad \text{for any timelike } t,$$

where G is the Einstein tensor and 1 is the metric. To take into account the rôle played by local conditions I introduce the notion of ‘C-spacetime’.

3. Definition. A smooth connected paracompact Hausdorff orientable manifold endowed with a Lorentzian metric is a *C-spacetime* if it satisfies a local condition C .

Correspondingly,

4. Definition. A C-spacetime M' is a *C-extension* of a C-spacetime M if M is isometric to an open proper subset of M' . M is called *C-extendible* if it has a C-extension and *C-maximal* otherwise.

Throughout the paper the letter C denotes the *same* local condition. I shall not specify it though, because it is important that all the results below are valid for *any* C including trivial. Obviously, in this latter case (when C is trivial, that is when, in fact, no additional local condition is imposed on the metric) ‘C-spacetime’ and ‘C-extension’ are the same as ‘spacetime’ and ‘extension’, respectively.

[‡] For other proposed criteria see [4].

Now we are in position to refine item (ii) in the above-formulated criterion: the words ‘allowed maximal extension’ stand there for ‘C-maximal C-extension’.

Imposing a local condition we still cannot provide uniqueness of evolution of a spacetime. Whenever M has a C-maximal C-extension M' (except when $M' = \overline{M}$) it has infinitely many other such extensions [5, 6] including, for example, those obtained from M' in the way we built the DP space from the Minkowski plane. I interpret this as impossibility of creating a prescribed spacetime: whatever initial data (i. e. the geometry of the ‘initial region’ M) are prepared, and whatever are the equations of motion of the matter filling M , one does not know whether M will evolve in the desired M' , or in any of other possible extensions satisfying the same local conditions. However, in building a time machine it does not matter *how exactly* the spacetime will evolve[§], but only whether *a* CTC will appear. So it would suffice to create a situation in which a CTC is present in *any* of the possible extensions (as is sometimes the case with singularities [7]: fulfillment of some local conditions guarantees the existence of a singularity in the extensions of some spacetimes even without fixing uniquely their evolution). On the other hand, (ii) is also a *necessary* condition for considering a CTC as artificial. In a theory (quantum gravity?) where different *probabilities* could be ascribed to extensions, it would not be the case. One could take credit for creating ℓ even if one’s activity did not lead to its inevitable appearance, but just increased its probability. However, any further discussion of this hypothetical theory is beyond the scope of the present paper.

Theorem. Any C-spacetime U has a C-maximal C-extension $M^{\max}$ such that all closed causal curves in $M^{\max}$ (if they exist there) are confined to the chronological past of U .

Summing up we can say, that the theorem does not exclude at all the possibility of a closed causal curve. But it shows that in manufacturing such a curve it does not matter whether one, say, moves the mouths of a wormhole, or just utters: ‘Abracadabra!’. The results will be exactly the same: the curve may appear and it may not as well.

5. Example. Consider a cylinder $C \equiv \{t \in \mathbb{R}^1, \psi = \psi + 2\pi\}$ with the metric 1: $ds^2 = -2d\psi dt - t d\psi^2$. The part $U \equiv \{p \in C: t(p) < 0\}$ of this cylinder is a causal spacetime called the Misner space [7], while the region $t \geq 0$ contains causal loops. To see whether these loops are an inevitable consequence of something that takes place in the Misner space consider all possible maximal extensions of the latter. Some of the extensions [e. g. $(C, 1)$] are acausal and some are not. Pick, for example, a function Ω such that

$$\Omega|_U = 1, \quad \Omega(p) = 0 \quad \Leftrightarrow \quad p \in \zeta,$$

where ζ is the ray $\psi = 0, t \geq 0$ (see figure 1b). The spacetime $(C', \Omega^{-1}1)$, where $C' \equiv C - \zeta$, is a causal and, most likely, maximal extension of U . However, not all of these extensions are equally relevant. As is discussed above, we can impose a local condition C and declare all spacetimes that do not obey it ‘unphysical’. For example,

[§] Actually, the fact that one cannot predict the evolution of a given spacetime is beneficial in building a time machine being the protection against the time travel paradoxes [6].

we could require that a spacetime should be flat (U is flat). Then $(C, 1)$ would be an ‘allowed’ extension (a C -extension) of U , while $(C', \Omega^{-1}1)$ would be not. What the theorem asserts in application to this case is: whatever C is chosen — as long as $(C, 1)$ obeys it — there can be found a *causal* maximal extension $M^{\max}$ of U also obeying C . Let, for example, $\{M_n\}$, $n = \dots, -1, 0, 1 \dots$ be a set of spacetimes each isometric to $(C', 1)$, i. e. M_n are flat cylinders with the vertical cuts. Then the spacetime obtained by gluing for each n the left bank of the cut in M_n to the right bank of the cut in M_{n+1} will be just a desired $M^{\max}$. That it satisfies any C obeyed by $(C, 1)$ follows from the fact that $M^{\max}$ is locally isometric to $(C, 1)$.

1.1. Warning!

From the next subsection on to make the text readable I omit the letter C and write just ‘spacetime’, ‘extension’, etc. instead of ‘ C -spacetime’, ‘ C -extension’, etc. This definitely is a misuse of terms, but perhaps not that awful because:

1. No confusion must arise, since *nowhere* below these words are used in their ‘usual’ sense;
2. As has already been mentioned C is not specified. It may, in particular, be trivial. Which means that all that is below remains true even if understood ‘literally’, i. e. if this warning is ignored. The only problem is that what would be proven in such a case is not the theorem formulated above, but only its weaker version (obtained from the original one by omitting C ’s);
3. It is quite easy to ‘restore the real meaning’ of any sentence below. It suffices to add a C to *each* of the words ‘spacetime’, ‘extension’, etc;
4. One need not keep this warning in mind all the time. If something is valid for *arbitrary* spacetimes, the fact that it is also valid for C -spacetimes is absolutely trivial in most cases. The only exception is the matters of existence and membership. Of course ‘ A is a spacetime’ does not necessarily imply ‘ A is a C -spacetime’. We shall encounter such not-absolutely-trivial situation only once — in proposition 40 — and shall take care to show explicitly that the relevant spacetime is a C -spacetime indeed.

1.2. Outline of the proof

Consider an extendible spacetime M . Let us first try to find an extension M_Δ (the reason for such notation becomes evident later) of M such that all causal loops, if they exist in M_Δ , lie in M . If we find such an M_Δ and if, in addition, it is maximal, the theorem would be proven.

Take an *arbitrary* extension M^{ext} of M and consider its open subset $W \equiv M \cup N$, where N is a *diamond* neighbourhood of a point p in the boundary of M . The precise meaning of the term ‘diamond’ (given in definition 15) is immaterial at the moment. It is important only that a diamond neighbourhood exists for any point (see proposition 16) and that diamond neighbourhoods are normal [8]. Clearly W is an extension of M . Moreover, N (being normal) does not contain any causal loops. Hence new (i. e. not

confined to M) causal loops may exist in W only if there is a causal curve through a point of $N - M$ such that both its endpoints lie in M (see the curve λ in figure 2). Which means that if we are lucky enough and $M_N \equiv M \cap N$ is *causally convex in N*




Figure 2. If we choose $N = D_3$, M_N will be not causally convex. But we can ‘unglue’ its lower component from M (so that it is not a part of M any longer and the dark region in the picture must be viewed as a part of M seen through N). $M \cup N$ in such a case have no new causal loops.

(i. e. any causal curve in N lies in M_N if both its ends do), then we can be assured that W is just a desired extension M_Δ . This condition can be slightly weakened. Instead of the causal convexity of M_N we can require that only a connected component $M_\vee$ of M_N is causally convex in N . To obtain M_Δ in such a case one only need ‘unglue’ all other components of M_N from M (see figure 2).

Generally, $M_\vee$ of course need not be causally convex. If, for example, M^{ext} is a cylinder $C \equiv \{x \in \mathbb{R}^1, t = t + 1\}$ with the metric $ds^2 = dx^2 - dt^2$ and M is the region bounded by the bold line in figure 2, then M_N is causally convex when $N = D_4$, and is not when $N = D_{1,2}$ (when $N = D_3$, there are two connected components in M_N , of which the upper one is causally convex and the lower is not). There are spacetimes, however, — I shall call them *locally causally convex*, or LCC — such that $M_\vee$ is causally convex whatever M^{ext} and M_N and whichever component of M_N are chosen (an example, as can be seen from proposition 19, is the Minkowski half-space $t < 0$). It follows from the above reasoning that any extendible LCC spacetime M has an extension M_Δ such that all causal loops in M_Δ are confined to M (cf. proposition 22).

Our next step is constructing in section 3 yet another extension of an LCC spacetime M . This new extension — denoted by $M_\mathcal{N}$ — is made from M_Δ by some cutting and gluing and possesses the following properties (it is the proof in sections 4,5 of the last two of them that constitutes the most technical and tiresome part of the whole proof):

- (i) It is locally isometric to M_Δ , which in its turn is a part of M^{ext} . That is how we know that C holds in $M_\mathcal{N}$ (which, thus, is a spacetime indeed, see the previous subsection);
- (ii) Like M_Δ it has no causal loops other than those lying in M ;
- (iii) Unlike M_Δ it is always locally causally convex if such was M .

Now that we see that any (extendible) LCC spacetime M can be extended to a larger (and *also* LCC) spacetime without the appearance of new causal loops, the theorem

can be proven by just employing the Zorn lemma as soon as we show that any U has an LCC extension M which contains no causal loops outside $I_M^-(U)$. This is done as follows. Consider (for a given U) the set $\mathbf{V}$ of all possible spacetimes of the form $I_{U'}^-(U)$. Clearly, all causal loops in any $V \in \mathbf{V}$ lie in the chronology past of U (just because it is the whole spacetime). So, all we need is to find an LCC element in $\mathbf{V}$. To this end we show (again by using the Zorn lemma) that there is a maximal element V^m in $\mathbf{V}$, i. e. such an element that no $V \in \mathbf{V}$ is an extension of V^m . In no extension of V^m can a past directed causal curve leave V^m . Whence V^m is LCC.

1.3. Notation and conventions

In this paper the signature is chosen to be $(-, +, +, +)$. Whenever possible I use capital Latin letters of different fonts to denote 4- and 3-dimensional sets (U , M , etc. for the former and $\mathcal{B}$, $\mathcal{S}$, etc. for the latter), and Greek capital letters to denote 2- and 1-dimensional sets of points. Small Greek and Latin letters will denote, as a rule, curves (and sometimes isometries) and points, respectively. Also the following notation will be used:

$$\langle p, r \rangle_U \equiv I_U^+(p) \cap I_U^-(r), \quad \leqslant p, r \geqslant_U \equiv J_U^+(p) \cap J_U^-(r), \quad \leqslant p, r \geqslant \equiv \leqslant p, r \geqslant_M,$$

where U is an open subset of a spacetime M .

6. Remark. If $a, b \in \langle x, y \rangle_U$, then obviously

$$\langle a, b \rangle_{\langle x, y \rangle_U} = \langle a, b \rangle_U \quad \text{and} \quad \leqslant a, b \geqslant_{\langle x, y \rangle_U} = \leqslant a, b \geqslant_U.$$

2. Various types of sets

In this section I introduce the notions of ‘diamond’ and ‘locally causally convex’ sets and for later use establish some basic properties of such sets. Some of the material of the section (definitions 7, 9, 10 and — most likely — propositions 8, 11) can be found elsewhere and is included to make the paper self-contained.

2.1. (Causally) convex sets

7. Definition. An open set O is *convex* if it is a normal neighbourhood of each of its points.

With any two points x, y a convex set O contains also a (unique) geodesic segment λ_{xy} that connects them. To an extent this property is shared by the closure of a convex set.

8. Proposition. If O is a convex subset of a convex spacetime M , $a \in O$ and $c, d \in \overline{O}$, then

$$\lambda_{ac} - c \subset O, \quad \lambda_{cd} \subset \overline{O}.$$

9. Definition. An open set O is called a *causally convex subset* of U if with any two points a, b it contains also the set $\langle a, b \rangle_U$.

Note that in contrast to convexity, causal convexity is not an intrinsic property of a set. That is, if U_1 is a convex and causally convex subset of M , then any $U_2 \subset M$ isometric to U_1 is also convex, but not necessarily causally convex.

2.2. Simple sets

10. Definition. An open set O is *simple* if it is convex and its closure is a compact subset of some other convex neighbourhood.

The Whitehead theorem ensures the existence of a simple neighbourhood of any point of any spacetime. Since a neighbourhood of a point is itself a spacetime (and since any its simple subset is at the same time a simple subset of the larger spacetime), this means that any point has ‘arbitrarily small’ simple neighbourhoods, or in other words that the simple neighbourhoods constitute a base of topology in any spacetime.

11. Proposition. Let O be convex. Then any connected component of $O \cap O'$ is convex (simple), if so does O' . Also if O is simple and its subsets O_1, O_2 are convex, then the following sets

$$O_1, \quad O_2, \quad O_1 \cap O_2$$

all are simple.

Simple sets are still ‘not simple enough’ for our needs. The problem is that there is no direct relation between simplicity and causal convexity. A timelike curve (provided it is not geodesic) leaving a simple set still can return in it. Below we shall overcome this problem by distinguishing a special subclass of convex sets. In doing so we shall lean upon the following fact.

12. Proposition. Any point q of any spacetime has a simple neighbourhood O such that the sets $I_O^\pm(p)$ are simple for any $p \in O$.

Proof. Let $\{e_{(i)}\}$ with $e_{(i)}^\mu e_{(j)\mu} = \eta_{ij}$ be a smooth frame field in some simple neighbourhood O' of q . Emitting all possible geodesics λ from each point of O' we introduce a normal coordinate system $X_{\{p\}}^\mu$ for each $p \in O'$ by the following procedure: for any point r we find the geodesic $\lambda_{pr}(\xi)$, where ξ is an affine parameter such that $p = \lambda_{pr}(0)$, $r = \lambda_{pr}(1)$, and ascribe to r the coordinates $X_{\{p\}}^\mu(r)$ equal to the coordinates of $\partial_\xi(p)$ in the basis $\{e_{(i)}(p)\}$. The functions $X_{\{p\}}^\mu(r)$ depend smoothly on both r and p . Hence, in particular, for any δ there exists a simple neighbourhood O_δ of q such that

$$\overline{O_\delta} \subset O', \quad \sup_{p, r \in O_\delta} X_{\{p\}}^\mu(r) < \delta, \quad \forall \mu. \quad (1)$$

We choose O_δ with sufficiently small δ to be the desired O . So to prove the proposition we only need to show that

$$\exists \delta: \quad \lambda_{ab} \subset I_{O_\delta}^\pm(p) \quad \forall p, a, b: \quad p \in O_\delta, \quad a, b \in I_{O_\delta}^\pm(p) \quad (2)$$

(strictly speaking (2) means that $I_O^\pm(p)$ are convex, but proposition 11 ensures that they are simple as well).

For causal λ 's (2) follows just from the definition of $I_{O_\delta}^\pm(p)$, so we can restrict ourselves to the spacelike ones:

$$l^\mu l_\mu = 1, \quad \text{where } l_\mu \equiv (\partial_\tau)_\mu. \quad (3)$$

Here τ is understood to be an affine parameter. Let us introduce the function

$$\sigma(r) \equiv 1(\mathbf{x}(r), \mathbf{x}(r)),$$

where $\mathbf{x}(r) \in T_r$ is the ‘position vector’ [8] defined for a fixed p by $x^\mu(r) = X_{\{p\}}^\mu(r)$. Since σ is a smooth function negative inside $I_{O_\delta}^\pm(p)$ and positive outside, (2) will be proved once we prove that, when δ is small enough, a spacelike geodesic can touch a null cone only from outside:

$$\sigma''(\tau_0) > 0, \quad (4)$$

$$\text{where } \sigma(\tau) \equiv \sigma \circ \lambda_{ab}(\tau), \quad ' \equiv \frac{d}{d\tau}, \quad \tau_0: \sigma(\tau_0) = 0, \quad \sigma'(\tau_0) = 0.$$

To obtain (4) let us first use the relation $\sigma_{,\mu} = 2x_\mu$ (proved e. g. in [8])

$$\sigma' = \sigma_{,\mu} l^\mu = 2x_\mu l^\mu. \quad (5)$$

This gives

$$\sigma'' = (2x_\nu l^\nu)_{,\mu} l^\mu = 2l^\nu l^\mu x_{\nu;\mu} \quad (6)$$

(in the last equation we used the fact that $\lambda_{ab}(\tau)$ is a geodesic). Now consider the point $\lambda_{ab}(\tau_0)$, where λ_{ab} touches the null cone, and for $\mathbf{z} \in T_{\lambda_{ab}(\tau_0)}$ denote by $T_{\mathbf{z}}^\perp$ the 3-dimensional subspace of $T_{\lambda_{ab}(\tau_0)}$ orthogonal to $\mathbf{z}$. It follows from (5) that $\mathbf{l}(\tau_0) \in T_{\mathbf{x}}^\perp$. Also $\mathbf{x} \in T_{\mathbf{x}}^\perp$, since $\mathbf{x}$ is null (recall that $\sigma(\tau_0) = 0$). Take a basis $\{\mathbf{x}, \mathbf{d}_{(1)}, \mathbf{d}_{(2)}\}$ in $T_{\mathbf{x}}^\perp$ defined (non-uniquely, of course) by the following relations:

$$\mathbf{d}_{(\alpha)} \in T_{\mathbf{x}}^\perp \cap T_{e(0)}^\perp, \quad 1(\mathbf{d}_{(\alpha)}, \mathbf{d}_{(\beta)}) = \delta_{\alpha\beta} \quad \alpha, \beta = 1, 2. \quad (7)$$

Decomposing $\mathbf{l}$ we see from (3) that

$$l^{(1)2} + l^{(2)2} = 1, \quad \text{where } \mathbf{l} = l^{(1)}\mathbf{d}_{(1)} + l^{(2)}\mathbf{d}_{(2)} + l^{(3)}\mathbf{x}. \quad (8)$$

The term $l^{(3)}\mathbf{x}$ in $\mathbf{l}$ gives no contribution to the right hand side of (6) because $x^\nu x_{\nu;\mu} = \sigma_{,\mu}/2 = x_\mu$ and $x^\mu x_{\nu;\mu}$ is proportional to x_ν (since $\mathbf{x}$ is tangent to a geodesic). Hence

$$\sigma'' = 2l^{(\alpha)}l^{(\beta)}d_{(\alpha)}^\nu d_{(\beta)}^\mu x_{\nu;\mu} = 2l^{(\alpha)}l^{(\beta)}d_{(\alpha)}^\nu d_{(\beta)}^\mu (g_{\mu\nu} + \Gamma_{\nu,\mu\rho}x^\rho). \quad (9)$$

With the notation

$$\Gamma \equiv 2|l^{(\alpha)}l^{(\beta)}d_{(\alpha)}^\nu d_{(\beta)}^\mu \sup_{\substack{\nu,\mu,\rho \\ r,p \in O'}} \Gamma_{\nu,\mu\rho}|$$

we find from (1,7,8,9) that

$$|\sigma'' - 2| \leq 4\Gamma\delta.$$

$l^{(\alpha)}$ are bounded [see (8)] and so obviously are $d_{(\alpha)}^\nu$. Thus Γ is finite and hence σ'' is positive for sufficiently small δ , which proves (4) and thereby the whole proposition. $\square$

2.3. IGH neighbourhoods

A few useful characteristics of a set are obtained by simply forgetting about the ambient space.

13. Definition. Let U be an open subset of a spacetime (M, g) . We call U *intrinsically (strongly) causal* if $(U, g|_U)$ is a (strongly) causal spacetime and *intrinsically globally hyperbolic* (IGH) if $(U, g|_U)$ is globally hyperbolic.

14. Proposition. Any point q of any simple set O has an IGH neighbourhood U of the form $\langle p, r \rangle_O$, where $p, r \in O$.

Proof. Consider two sequences $p_m, r_m \in O$:

$$p_m, r_m \rightarrow q, \quad q \in Q_m \equiv \leq p_m, r_m \geq_O.$$

Each set Q_m is closed in the topology of O [7, prop. 4.5.1], but it well may be not closed in the topology of the ambient spacetime M . When the latter is true there exists a point $x_m \in \overline{Q_m} \cap \text{Bd } O$. If x_m would exist for infinitely many m , there would be a subsequence x_n converging to some $x \in \text{Bd } O$ and the sequences of geodesics $\lambda_{p_n x_n}$ and $\lambda_{r_n x_n}$ (of which the former are future- and the latter are past-directed) would converge to the same geodesic λ_{qx} , which thus would be both future- and past-directed at once. This is impossible and hence there exists m_0 such that Q_{m_0} is closed in the topology of M , or in other words (recall that $\overline{O}$ is compact)

$$\exists m_0 : \quad Q_{m_0} = \leq p, r \geq_O \quad \text{is compact,} \quad \text{where } p \equiv p_{m_0}, \quad r \equiv r_{m_0}.$$

Being a subset of the simple neighbourhood O the set $U \equiv \langle p, r \rangle_O$ is intrinsically strongly causal [9, prop. 4.10]. So, to prove that it is IGH it remains only to show that $\leq a, b \geq_U$ is compact for any $a, b \in U$. Which follows from the fact (see remark 6) that

$$\leq a, b \geq_U = \leq a, b \geq_O$$

and the right hand side is a closed subset of the compact Q_{m_0} .

□

2.4. Diamond sets

15. Definition. Let $R(D)$ be a set consisting of a spacetime D and all its subsets of the form $\langle x, y \rangle$. We call D *diamond*[†] if any $A \in R(D)$:

(i) is convex; (ii) is IGH; and (iii) with any two points a, b contains also points c, d such that $a, b \in \langle c, d \rangle_A$.

16. Remark. It follows from remark 6 that $A \in R(D)$ implies $R(A) \subset R(D)$. Therefore, if D is IGH or diamond then so does any $A \in R(D)$.

[†] The name refers to the shape of an obvious diamond spacetime — the region $\langle a, b \rangle$, where a and b are some points in the Minkowski plane. After this paper had been written I recalled that Yurtsever already used the term ‘diamond’ to denote another type of sets [10]. The contexts however are so different that no confusion must arise.

With so many good qualities diamond sets are, in the Lorentzian case, a good candidate for the role fulfilled in the Riemannian case by balls. The following proposition shows that they also constitute a base of topology of spacetime.

17. Proposition. Any point q of any spacetime has a diamond neighbourhood.

Proof. Let O and $U \subset O$ be the neighbourhoods of q from propositions 12 and 14, respectively. Then any $A \in R(U)$ is, first, simple (by propositions 11,12) and, second IGH (by proposition 14 coupled with remark 16). Condition (iii) obviously also holds in A . So U can be taken as the desired neighbourhood. $\square$

2.5. Locally causally convex sets

Now we are in position to introduce the notion that plays the central part in our proof — local causal convexity, which is an analog of causal convexity, but in contrast to the latter characterizes the set itself and does not depend on the way it is embedded into a larger space. Let D be a diamond subset of an extension M^e of a spacetime M , and $D_\vee$ be a connected component of $D \cap M$.

18. Definition. M is *locally causally convex* (LCC) if for any M^e and D each $D_\vee$ is a causally convex subset of D .

Generally, neither convex, nor IGH sets are LCC (consider a rectangle in the Minkowski plane, and the ‘bad’ set from [11], respectively). However the following holds.

19. Proposition. Any IGH convex spacetime M is LCC.

Proof. Consider a future-directed timelike curve $\gamma \subset D$ from a to b , where $a, b \in D_\vee$. We must show that the whole γ lies in $D_\vee$.

Let $c \in \gamma \cap D_\vee$. By proposition 11 $D_\vee$ is convex. Therefore a geodesic $\lambda_{ac} \subset D_\vee$ from a to c must exist. λ_{ac} is future-directed [D is convex, so λ_{ac} is the unique geodesic connecting a and c in D , while the existence of γ guarantees that $c \in I_D^+(a)$] and thus $c \in \gamma \cap J_{D_\vee}^+(a)$. Similarly, $c \in \gamma \cap J_{D_\vee}^-(b)$, and so,

$$\gamma \cap D_\vee = \gamma \cap \leq a, b \geq_{D_\vee}.$$

The right hand side is compact (since both M and D are IGH and convex), and $D_\vee$ is open. Consequently, $\gamma \subset D_\vee$. $\square$

20. Corollary. Any diamond spacetime is LCC (and, accordingly, any point has an arbitrarily small intrinsically causal LCC neighbourhood).

The reverse, of course, is not true. One of the reasons is that local causal convexity is, loosely speaking, a characteristic of the ‘superficial’ (i. e. lying ‘near the boundary’) regions of a spacetime rather than its bulk. This, in particular, entails quite regular structure of the boundary of an LCC region: ‘mostly’ it is achronal (though not *always* by the reasons obvious from inspection of D_4 in figure 2).




Figure 3. Of the two curves meeting in p the upper one is $\mathcal{B}^+$ and the lower one is $\mathcal{B}^-$. p does not belong to either.

21. Proposition. If M is an LCC subset of a spacetime M_1 and U is a neighbourhood of some point of $\text{Bd } M$, then within U there always exists a diamond set H such that, $\text{Bd } H_v \cap H$ (where H_v is a connected component of $H \cap M$) is a closed, imbedded, achronal three-dimensional C^{1-} submanifold in H .

Proof. We begin by proving that there is a timelike curve in M whose end point lies in $\text{Bd}_U M$ (the boundary of M in U). Let U_M be a connected component of $U \cap M$ and let $p \in \text{Bd}_U U_M$ (see figure 3). Denote by $\mathcal{B}^\pm$ the (maybe empty) sets of points of $\text{Bd}_U U_M$ which are the future (past) endpoints of timelike curves lying in U_M .

Clearly, any timelike curve in U connecting a point $a \in U_M$ with p contains at least one point of $\mathcal{B}^+ \cup \mathcal{B}^-$. In particular, if $\mathcal{B}^+ = \emptyset$, then $I_U^-(p) \subset U - U_M$. And hence if also $\mathcal{B}^- = \emptyset$, then $I_U^+(I_U^-(p)) \subset U - U_M$, which is impossible since $I_U^+(I_U^-(p))$ is a neighbourhood of p . So

$$\mathcal{B}^- \cup \mathcal{B}^+ \neq \emptyset.$$

Suppose for definiteness that it is $\mathcal{B}^+$ that is non-empty:

$$\mathcal{B}^+ \neq \emptyset.$$

Our next step is to show that $\mathcal{B}^+$ and $\mathcal{B}^-$ are separated. Let $q \in \mathcal{B}^+$. By the definition of $\mathcal{B}^+$, q lies in $I_U^+(U_M)$ and hence so does some its neighbourhood Q . Without loss of generality (see proposition 17) U can be taken diamond. Then $q \notin \overline{\mathcal{B}^-}$ since otherwise some points of $\mathcal{B}^-$ also would lie in Q . Such points would belong to $X \equiv (I_U^+(U_M) \cap I_U^-(U_M)) - U_M$, but M is LCC, U is diamond, and so, X must be empty by definition 18. Thus

$$\mathcal{B}^\pm \cap \overline{\mathcal{B}^\mp} = \emptyset. \tag{10}$$

It follows that for a sufficiently small neighbourhood $H \subset U$ of q

$$H \cap \mathcal{B}^- = \emptyset. \quad (11)$$

To complete the proof it suffices now to require that H be diamond (which by proposition 17 is always possible) and to choose as $H_\vee$ a component of $H \cap M$ lying in U_M . Any past directed timelike curve in H leaving $H_\vee$ (and thus also U_M) would contain a point of $\mathcal{B}^-$, which is impossible by (11). So $H_\vee$ is the past set in H and hence by [7, proposition 6.3.1] its boundary is a closed, imbedded, achronal three-dimensional C^{1-} submanifold of H .

□

3. Construction of $M_\mathcal{N}$

In this section for an arbitrary extendible LCC spacetime M we construct an extension $M_\mathcal{N}$ of a special type (as will be proved in the subsequent sections $M_\mathcal{N}$ is LCC and has no closed causal curves except those lying in M). $M_\mathcal{N}$ will be built in a few steps. First we glue a diamond region H to M obtaining thus an extension M_Δ (see figure 3b). Then to the ‘upper’ (that is lying outside M) part of M_Δ we glue yet another copy of H (in doing so we remove a three-dimensional surface, so that the resulting spacetime $M_\Diamond$ (depicted in figure 4a) be Hausdorff). Finally, a smaller diamond set H' is glued to $M_\Diamond$ (see figure 5).

3.1. The spacetime M_Δ .

22. Proposition. Any extendible LCC spacetime M has an extension M_Δ such that:

- (I). $M_\Delta = M \cup H$, where H is diamond, and $M \cap H$ is connected;
- (II). M is a past set in M_Δ , or (see remark 23 below)
- (II'). M is a future set in M_Δ .

Proof. Let M_1 , H , and $H_\vee$ be as in proposition 21. Let further $\overset{*}{M}$ and $\overset{*}{H}$ be spaces isometric to M and H , respectively:

$$\phi_M: M \rightarrow \overset{*}{M}, \quad \phi_H: H \rightarrow \overset{*}{H}, \quad \phi_M, \phi_H \text{ — isometries.}$$

The spacetime $M_2 \subset M_1$ defined by $M_2 \equiv M \cup H$ can be presented as a result of ‘gluing’ $\overset{*}{H}$ to $\overset{*}{M}$ by an isometry:

$$M_2 = \overset{*}{M} \cup_\phi \overset{*}{H},$$

where

$$\phi \equiv \phi_M \circ \phi_H^{-1}: \phi_H(H \cap M) \rightarrow \phi_M(H \cap M).$$

We construct M_Δ by ‘ungluing’ $\overset{*}{H}$ from $\overset{*}{M}$ along all but $H_\vee$ connected components of $H \cap M$ (see figure 3):

$$M_\Delta \equiv \overset{*}{M} \cup_{\phi|_{\phi_H(H_\vee)}} \overset{*}{H}.$$




Figure 4. M_Δ is obtained by gluing $\tilde{K} = \tilde{H} - \overline{\tilde{S}_-}$ to $M_\Delta - \overline{\tilde{S}_0}$.

From now on by M , H , and $H_\vee$ we understand the corresponding regions of M_Δ (this must not lead to any confusion, since we shall not consider M_1 any more). M_Δ is locally isometric to a part of M_1 and hence C holds in it. So, it is a C-extension of M , indeed. Further, M_1 satisfies (I) by construction and (II), or (II') (depending on whether q was taken in $\mathcal{B}^+$, or in $\mathcal{B}^-$) by the reasons discussed in proposition 21. □

23. Remark. In what follows we assume for definiteness that it is (II) that holds for our M_Δ .

24. Remark. Condition (II) of course implies that the boundary $\mathcal{S} \equiv \text{Bd } M$ is a closed, imbedded, achronal three-dimensional C^{1-} submanifold in M_Δ . This surface divides H into two parts: $H_\vee$ and $H_\wedge \equiv H - \overline{H_\vee}$.

25. Remark. By the definition of local causal convexity condition (I) implies that all causal loops in M_Δ (if there are any) are confined to M . If in addition M_Δ always were LCC, which unfortunately is not the case, we would not need anything below up to proposition 40.

3.2. The spacetime M_Δ .

Now we want to construct for M yet another extension, which we shall denote by $M_\Diamond$ (and which is not an extension of M_Δ). We shall do this similarly to the way we constructed M_Δ that is by, first, presenting some auxiliary spacetime ($M_\Delta - \Theta$, see below) as a result of gluing together two spacetimes ($\tilde{K}$ and $\hat{M}_{\bowtie}$) and by then ungluing them along a connected component of their intersection.

Let $H' \subset H$ be a diamond neighbourhood with the compact closure in H and let it intersect $\mathcal{S}$, thus splitting the latter into three non-empty disjoint parts (see figure 4):

$$\mathcal{S}_- \equiv \mathcal{S} \cap H', \quad \mathcal{S}_0 \equiv \mathcal{S} - \overline{\mathcal{S}_-}, \quad \Theta \equiv \overline{\mathcal{S}_-} - \mathcal{S}_-.$$

$\mathcal{S}_-$ in its turn divides H' into two disjoint regions:

$$H'_\vee \equiv H' \cap H_\vee, \quad H'_\wedge \equiv H' \cap H_\wedge.$$

Now let $\hat{M}_\Delta$ and $\tilde{H}$ be spaces isometric to M_Δ and H , respectively. The isometries are:

$$\psi_M: M_\Delta \rightarrow \hat{M}_\Delta, \quad \psi_H: H \rightarrow \tilde{H}.$$

We shall often write $\tilde{A}$ for $\psi_H(A)$ and sometimes $\hat{A}$ for $\psi_M(A)$. In particular:

$$\tilde{H}_\wedge \equiv \psi_H(H_\wedge), \quad \tilde{H}'_\vee \equiv \psi_H(H'_\vee), \quad \tilde{\mathcal{S}}_- \equiv \psi_H(\mathcal{S}_-), \quad \hat{\mathcal{S}}_- \equiv \psi_M(\mathcal{S}_-), \text{ etc.}$$

Consider the spacetime $M_\Delta - \Theta$. Obviously,

$$M_\Delta - \Theta = \hat{M}_\Delta \cup_{\psi_{MH}} \tilde{K},$$

where

$$M_\Delta \equiv M_\Delta - \overline{\mathcal{S}_0}, \quad K \equiv H - \overline{\mathcal{S}_-}, \quad \psi_{MH} \equiv \psi_M \circ \psi_H^{-1}.$$

Note that $M_\Delta \cap K$ consists of two disjoint regions: $H_\vee$ and $H_\wedge$, which enables us to build a new spacetime by ungluing K from M_Δ along one of them. Namely, we define

$$M_\diamond \equiv \hat{M}_\Delta \cup_\psi \tilde{K}, \quad \text{where } \psi \equiv \psi_{MH}|_{\tilde{H}_\wedge}. \quad (12)$$

The definition (12) produces two natural isometries:

$$\varpi_{M_\Delta}: \hat{M}_\Delta \rightarrow M_\diamond, \quad \varpi_K: \tilde{K} \rightarrow M_\diamond.$$

From now on for the subsets $\varpi_{M_\Delta}(\hat{M}_\Delta)$, $\varpi_{M_\Delta}(\tilde{K})$, $\varpi_{M_\Delta}(\hat{\mathcal{S}}_-)$, etc. of $M_\diamond$ we shall write[‡] simply M_Δ , K , $\mathcal{S}_-$, etc., while the images of ϖ_K we shall mark by $\diamond$, i. e. $\varpi_K(\tilde{K})$, $\varpi_K(\tilde{H}'_\vee)$, $\varpi_K(\tilde{H}_\vee)$, etc. we shall denote by $K_\diamond$, $H'_{\diamond\vee}$, $H_{\diamond\vee}$, etc. In this notation $M_\Delta \cap K_\diamond = H_\wedge$, for example.

3.3. The spacetime $M_\mathcal{N}$.

We shall be interested in one particular type of extensions of $M_\diamond$, which we obtain by pasting one more copy of H' to $M_\diamond$. More specifically, we take a spacetime $H'_\mathcal{N} = \varsigma(\tilde{H}'_\wedge)$ (where ς is an isometry) and define a new spacetime as follows

$$M_\mathcal{N} \equiv M_\diamond \cup_\chi H'_\mathcal{N}, \quad \text{where } \chi \equiv \varsigma \circ \varpi_K^{-1}|_{H'_{\diamond\vee}}.$$

It is easy to see that $M_\mathcal{N}$ is an extension of $M_\diamond$ [indeed, $M_\mathcal{N} - \overline{M_\diamond} = H'_{\mathcal{N}\wedge} \equiv \varsigma(\tilde{H}'_\wedge)$, so it is an extension of $M_\diamond$ in the ‘usual’ sense (see the introduction) and C holds in $M_\mathcal{N}$ since the latter is locally isometric to the corresponding part of M_1 , cf. proposition 22].

26. Remark. Consider the region $N \subset M_\mathcal{N}$ (depicted in figure 5) and the projection $\pi: N \rightarrow \tilde{H}$ defined as follows:

$$N \equiv M_\mathcal{N} - (M - H'_\vee) = \overline{H'_\vee} \cup K_\diamond \cup \overline{H'_{\mathcal{N}\wedge}},$$

$$\pi|_{H'} \equiv (\varpi_{M_\Delta} \circ \psi_{MH})^{-1}, \quad \pi|_{K_\diamond} \equiv \varpi_K^{-1}, \quad \pi|_{H'_\mathcal{N}} \equiv \varsigma^{-1}.$$

Clearly N is an area in the universal covering of $\tilde{H} - \tilde{\Theta}$.

[‡] Strictly speaking, this is some abuse of notation because originally we took, say, M_Δ to be a part of M_Δ (not of $M_\diamond$) but it must not lead to any confusion, since we shall not consider M_Δ any more.




Figure 5. M_N can be viewed as a result of gluing N to M .

4. The structure of M_N

Our proof in section 5 of that M_N is LCC will be based on the fact that in its extensions some curves with the same ends are nonhomotopic, and thus cannot belong to the same diamond set. What makes the curves nonhomotopic is a singularity that is present in M_N in spite of the fact that M_N was assembled of a few spacetimes each free from singularities. The nature of this singularity is precisely the same as in the Deutsch-Politzer spacetime, or, say, in the double covering of the Minkowski plane with a deleted point.

In this section we, first, establish that indeed there is a singularity in M_N and then consider, among other things, the ensuing restrictions on homotopic curves. Some of the facts concerning intersection of curves with the surfaces $S_{\pm}$ and $H'_{\wedge, \vee}$ seem obvious, but have to be *proved* because these surfaces are not sufficiently smooth to fall under the standard results.

4.1. The singularity

Let us present M_N in the following form:

$$M_N = M_{\bowtie} \cup K_{\diamond} \cup H'_N. \quad (13)$$

By construction (see proposition 22) M is a past set in $M_{\bowtie}$. So (since $K_{\diamond} \cup H'_N$ is open and has no common points with M) M is a past set in M_N as well. Similarly, $H'_{N\wedge}$ is a future set in H'_N and has no common points with $M_{\bowtie} \cup K_{\diamond}$ and hence is a future set in M_N (and likewise it can be shown that $H_{\wedge}$ and $H_{Q\vee}$ are, respectively, a future and a past sets in M_N). Which means that both $S_- = \text{Bd } M$ and $S_+ \equiv \text{Bd } H'_{N\wedge}$ are imbedded three-dimensional C^{1-} submanifolds achronal in M_N . The intersection of S_+ and S_- is empty, their union we shall denote by $S_{\bowtie}$. The following proposition, loosely speaking, says that Θ cannot be glued back into the spacetime.

27. Proposition. In any extension M^e of the spacetime M_N the sets $\mathcal{S}_\pm$ are closed.

Proof. It is easy to check that $\mathcal{S}_+$ is closed if so does $\mathcal{S}_-$. So we shall only prove the proposition for the latter. Suppose $\{a_n^{(1)}\}$ is a sequence of points such that (contrary to our claim):

$$a_n^{(1)} \in \mathcal{S}_-, \quad \exists \lim a_n^{(1)} = a \notin \mathcal{S}_-$$

and $\hat{a}_n^{(1)}$ are the images of these points in $\hat{M}_\Delta$:

$$\hat{a}_n^{(1)} \equiv \psi_M(a_n^{(1)}).$$

$\hat{a}_n^{(1)}$ are confined to the compact set $\overline{\psi_M(\mathcal{S}_-)}$ and therefore there exist points x , $\hat{q}$, and q :

$$x \equiv \lim \hat{a}_n^{(1)}, \quad \hat{q} \in I_{\hat{M}_\Delta}^+(x), \quad q \equiv \psi_M^{-1}(\hat{q}) \in M_N.$$

Obviously, $x \in \psi_M(\Theta)$ and so we can find a sequence $\{\hat{a}_n^{(2)}\}$:

$$\hat{a}_n^{(2)} \in \psi_M(\mathcal{S}_0), \quad \lim \hat{a}_n^{(2)} = x.$$

Now let $\{\hat{v}_n^{(i)}\}$ and $\{v_n^{(i)}\}$ with $i = 1, 2$ be the sequences in $T_{\hat{q}}$ and T_q , respectively, defined by

$$\hat{v}_n^{(i)} \equiv \exp_{\hat{q}}^{-1}(\hat{a}_n^{(i)}), \quad v_n^{(i)} \equiv d\psi_M^{-1}[\hat{v}_n^{(i)}].$$

Clearly,

$$\lim v_n^{(i)} = v, \quad \text{where } v \equiv d\psi_M^{-1}[\hat{v}], \quad \hat{v} \equiv \lim \hat{v}_n^{(i)}.$$

It follows from

$$a = \lim[\exp_q(v_n^{(1)})] = \exp_q\{d\psi_M^{-1}[\lim d\psi_M[v_n^{(1)}]]\} = \exp_q(v)$$

that the existence of a implies the existence (for sufficiently small positive ϵ) of a point a' :

$$a' \equiv \exp_q[(1 + \epsilon)v] = \lim \exp_q[(1 + \epsilon)v_n^{(2)}].$$

Each point

$$\exp_q[(1 + \epsilon)v_n^{(2)}] = \psi_M^{-1}(\exp_{\hat{q}}[(1 + \epsilon)\hat{v}_n^{(2)}])$$

lies in M (recall that $\hat{v}_n^{(2)}$ is timelike and $\hat{M}$ is a past set in $\hat{M}_\Delta$) and hence $a' \in \overline{M}$. Moreover, since v also is timelike

$$a' \in M.$$

But repeating the same reasoning for ψ_H and $\tilde{a}_n^{(i)} \equiv \psi_H \circ \psi_M^{-1}(\hat{a}_n^{(i)})$ instead of ψ_M and $\hat{a}_n^{(i)}$, respectively, one finds that $a' \in K_\diamond$, which is impossible, because by construction $K_\diamond$ and M are disjoint.

□

4.2. Curves intersecting $\mathcal{S}_\infty$

28. Definition. Let $\lambda: [0, 1] \rightarrow M^e$ be a timelike curve in an extension M^e of the spacetime M_N . We call τ_i a *positive (negative) root* if $\lambda(\tau_i) \in \mathcal{S}_\pm$. The number of the positive (negative) roots of λ we denote by $n_\pm[\lambda]$.

Obviously for any future-directed $\lambda \subset M_N$ the following holds:

$$\lambda(0) \in \overline{M}, \lambda(1) \notin M \quad \Rightarrow \quad n_-[\lambda] = 1, \quad \text{otherwise } n_-[\lambda] = 0 \quad (14a)$$

$$\lambda(0) \notin H'_{N\wedge}, \lambda(1) \in \overline{H'_{N\wedge}} \quad \Rightarrow \quad n_+[\lambda] = 1, \quad \text{otherwise } n_+[\lambda] = 0 \quad (14b)$$

Consider a homotopy $\lambda_\xi(\tau)$ with $\xi \in [0, 1]$ such that λ_ξ are timelike and future-directed. We shall denote the curves $\lambda_\xi|_{\tau=\tau_0}$ by $\mu_{\tau_0}(\xi)$.

29. Proposition. If the curves $\mu_{0,1}$ do not intersect $\mathcal{S}_\infty$, then $n_\pm[\lambda_0] = n_\pm[\lambda_1]$.

Proof. Suppose τ_0 is not a root of λ_{ξ_0} . Then the point $\lambda_{\xi_0}(\tau_0)$ does not belong to the closed set $\mathcal{S}_\infty$ and therefore some its neighbourhood U also does not intersect $\mathcal{S}_\infty$. So, around the point (τ_0, ξ_0) there exists a rectangle

$$\square_{\delta\epsilon} \equiv \{\tau, \xi: |\tau - \tau_0| < \delta, |\xi - \xi_0| < \epsilon\}$$

that does not contain roots.

Now suppose $\tau_0 \neq 0, 1$ is a root (positive for definiteness) of λ_{ξ_0} . The surface $\mathcal{S}_+$ lies in the spacetime M_N , where it bounds the future set $H'_{N\wedge}$, hence for any (sufficiently small) δ

$$\lambda_{\xi_0}(\tau < \tau_0) \in M_N - \overline{H'_{N\wedge}}, \quad \lambda_{\xi_0}(\tau > \tau_0) \in H'_{N\wedge},$$

when $|\tau - \tau_0| < \delta$.

So, when ϵ_a and δ_a are sufficiently small, any segment $\lambda_\xi(|\tau - \tau_0| < \delta_a)$ with $|\xi - \xi_0| < \epsilon_a$ will also lie in M_N and have its ends one in $H'_{N\wedge}$ and the other outside. Which by (14) implies that in $\square_{\delta_a\epsilon_a}$ there is exactly one root for each ξ .

Thus, any point $a = (\tau_0, \xi_0)$ lies in the center of a rectangle such that the number of the roots of λ_ξ located between $\tau_0 - \delta_a$ and $\tau_0 + \delta_a$ does not change for ξ varying between $\xi_0 - \epsilon_a$ and $\xi_0 + \epsilon_a$. Hence (due to the compactness of λ_ξ) such a positive constant ϵ_{ξ_0} can be found for any ξ_0 that $n_\pm[\lambda_\xi]$ does not change for $\xi \in (\xi_0 - \epsilon_{\xi_0}, \xi_0 + \epsilon_{\xi_0})$. As ξ varies over a compact set $[0, 1]$, its whole range can be covered by a finite number of such intervals with constant $n_\pm[\lambda]$. □

30. Corollary. If $\tau_n = \tau_n(\xi)$ is the n th root of λ_ξ , then $\mu_{\tau_n}(\xi)$ is a continuous curve. $\mu_{\tau_n}(\xi)$ and $\mu_{\tau_k}(\xi)$ are disjoint, when $n \neq k$

31. Remark. All timelike curves lying in a convex neighbourhood and connecting the same two points have equal $n_\pm$. Thus, for a convex neighbourhood we can speak about $n_\pm[pq]$ understanding by it $n_\pm[\lambda_{pq}]$, where λ_{pq} is an arbitrary timelike curve lying in this neighbourhood and connecting the points p and q . We also set $n_\pm[pq] = 0$ for $p = q$.

32. Remark. If points a, b, q_1, q_2 lie in a convex spacetime, $a, b \notin \mathcal{S}_\infty$, and $a \succeq b \succeq q_1, q_2$ (by $x \succeq y$ we mean $x \succ y$, or $x = y$), then

$$n_\pm[aq_2] - n_\pm[aq_1] = n_\pm[bq_2] - n_\pm[bq_1]. \quad (15)$$

Let $D \subset M^e$ be a diamond set and D_M be a connected component of $D \cap M_\mathcal{N}$. The following two propositions will help us to use the fact that M and $H'_{\mathcal{N}^\wedge}$ are LCC in reducing the proof of the local causal convexity of $M_\mathcal{N}$ to examining $K_\diamond$.

33. Proposition. If a future-directed timelike curve $\lambda \subset D$ starting and ending in D_M does not intersect $\mathcal{S}_\infty$, then its ends lie either both in M , or both in $H'_{\mathcal{N}^\wedge}$, or both in $K_\diamond$.

Proof. Consider a curve $\varphi: [0, 1] \rightarrow D_M$ connecting the ends of λ . Cover φ by a finite number of diamond subsets F_n of D_M :

$$\varphi \subset \bigcup_{n=1, \dots, N} F_n \subset D_M, \quad \varphi(0) = \lambda(0) \in F_1, \quad \varphi(1) = \lambda(1) \in F_N$$

and pick $N + 1$ points $q_n \notin \mathcal{S}_\infty$ such that:

$$q_1 \equiv \lambda(0), \quad q_{N+1} \equiv \lambda(1), \quad q_n \in F_{n-1} \cap F_n \quad \text{for } n \neq 1, N + 1.$$

For any pair q_n, q_{n+1} (since they belong to the same diamond set F_n) we can find a point $b_n \in I_{D_M}^+(q_n) \cap I_{D_M}^+(q_{n+1}) - \mathcal{S}_\infty$.

Pick a point a such that

$$a \in I_D^+(b_n) - \mathcal{S}_\infty \quad \forall n$$

(it is always possible because D is diamond). Applying twice equation (15) we get:

$$\begin{aligned} 0 = n_\pm[\lambda] &= n_\pm[a, \lambda(0)] - n_\pm[a, \lambda(1)] \\ &= n_\pm[q_1, b_1] - n_\pm[b_1, q_2] + n_\pm[q_2, b_2] - \dots - n_\pm[b_N, q_{N+1}]. \end{aligned} \quad (16)$$

Combining this with (14) we prove the proposition. □

34. Proposition. The sets $D_M \cap M$, $D_M \cap K_\diamond$ and $D_M \cap H'_{\mathcal{N}^\wedge}$ are connected.

Before proving the proposition we have to establish a lemma. Let $\lambda_\xi(\tau)$ be a homotopy considered in proposition 29 with an additional requirement that λ_ξ for each ξ is a geodesic in D . We denote by $\varphi(\xi) \equiv \lambda_\xi(\tau'(\xi))$ a curve (lying in the surface $\lambda(\xi, \tau)$) defined by a continuous function $0 \leq \tau'(\xi) \leq 1$ and consider the segment $\bar{\lambda}_\xi$ of λ_ξ between φ and an intersection of λ_ξ with $\mathcal{S}_\infty$.

35. Lemma. If $\varphi(\xi) \subset M_\mathcal{N}$ and the condition

$$\bar{\lambda}_\xi \equiv \lambda_\xi([\tau_n, \tau']) \subset M_\mathcal{N} \quad (17)$$

holds for $\xi = 0$ then it holds for all the rest ξ as well.

Proof. We shall prove the lemma for a negative root τ_n (the case of a positive root can be handled in much the same way). Then (17) is equivalent (recall that $H_\wedge$ is a future set and M is a past set in M_N) to

$$\overset{\square}{\lambda}_\xi \subset U \equiv I_{H_\wedge}^+(\mathcal{S}_-) \cup \mathcal{S}_- \cup M. \quad (18)$$

Let $\Xi \subset [0, 1]$ be the set of all ξ for which (18) [and hence (17)] holds and let ξ_* be a limit point of this set:

$$\xi_* = \lim \xi_k, \quad \xi_k \in \Xi.$$

We want to show that $\overset{\square}{\lambda}_{\xi_*} \subset U$. For $\tau_n(\xi_*) = \tau'(\xi_*)$ it is trivial. For $\tau_n(\xi_*) > \tau'(\xi_*)$ (i. e. a past-directed $\overset{\square}{\lambda}_{\xi_*}$) it follows from the fact that both $\lambda_{\xi_*}(\tau_n - 0)$ and $\lambda_{\xi_*}(\tau')$ lie in a connected component of the intersection of LCC M and diamond D . Finally, consider the case $\tau_n(\xi_*) < \tau'(\xi_*)$ (a future-directed $\overset{\square}{\lambda}_{\xi_*}$). By assumption $\lambda_{\xi_*}(\tau') = \varphi(\xi_*) \in M_N$. On the other hand, it is a limit point of $\lambda_{\xi_k}(\tau')$, so it lies in $I_{H_\wedge}^+(\mathcal{S}_-) \cap M_N$ and thus in N . The points $\pi[\lambda_{\xi_*}(\tau_n)]$ and $\pi[\lambda_{\xi_*}(\tau')]$ can be connected in $\tilde{H}$ by a geodesic γ . Since $\tilde{H}$ and D are convex

$$\gamma = \lim \pi(\overset{\square}{\lambda}_{\xi_k}) = \pi(\lim \overset{\square}{\lambda}_{\xi_k}) = \pi(\overset{\square}{\lambda}_{\xi_*})$$

and therefore (recall that $\overset{\square}{\lambda}_{\xi_*}$ is timelike)

$$\gamma - \gamma(\tau_n) \subset I_{\tilde{H}_\wedge}^+(\tilde{\mathcal{S}}_-).$$

Consequently, $\overset{\square}{\lambda}_{\xi_*} = \pi^{-1}(\gamma)$ lies in $I_{H_\wedge}^+(\mathcal{S}_-) \cup \mathcal{S}_-$, i. e. in U . Which means that Ξ is closed. It is also non-empty and obviously open. So, $\Xi = [0, 1]$. □

Proof of proposition 34. Let $a, b \in D_M \cap M$ and let a curve $\varphi \subset D_M$ from a to b be such that for some $x \in D$

$$\varphi \subset I_D^+(x).$$

Consider a homotopy $\lambda_\xi(\tau)$ with $\lambda_\xi \subset D$ being a geodesic segment from x to $\varphi(\xi)$. Suppose $\varphi \not\subset M$ and $p = \lambda_{\xi_p}(\tau_i)$, $q = \lambda_{\xi_q}(\tau_j)$ are, respectively, the first and the last points of φ that do not lie in M , which means of course that τ_i and τ_j are (i th and j th) roots of λ_{ξ_p} and λ_{ξ_q} , respectively. By lemma 35 the whole segment $\lambda_{\xi_q}([\tau_i, \tau_j])$ belongs to M_N . But a timelike curve in M_N cannot intersect $\mathcal{S}_-$ more than once. So actually $j = i$.

It follows then from corollary 30 that there is a curve $\mu_{\tau_i} \subset \mathcal{S}_-$ from p to q consisting of points $\lambda_\xi(\tau)$ and thus lying in D_M . So, the curve composed of the segment of φ from a to p , μ_{τ_i} , and the segment of φ from q to b can by a small variation (sending all inner points of this curve slightly to the past) be transformed into a curve $\varphi' \subset D_M \cap M$.

The connectedness of $D_M \cap K_\diamond$ and $D_M \cap H'_{N_\wedge}$ is proved in the same manner. □

4.3. Geodesics intersecting the boundary of $\tilde{H}'$

Let us introduce two new sets (see figure 4b): $\mathcal{H}'_{\wedge} \equiv \text{Bd}_{\tilde{H}} \tilde{H}'_{\wedge} - \tilde{\mathcal{S}}_-$ and $\mathcal{H}'_{\vee} \equiv \text{Bd}_{\tilde{H}} \tilde{H}'_{\vee} - \tilde{\mathcal{S}}_-$ (note that generally the surfaces $\mathcal{H}'_{\wedge, \vee}$ may not be achronal in $\tilde{H}$). It is easy to see that $\mathcal{H}'_{\wedge, \vee}$ are compact and that

$$\mathcal{H}'_{\vee} \cap \mathcal{H}'_{\wedge} \subset \tilde{\Theta}. \quad (19)$$

Let $\lambda_{\xi}(\tau)$ with $\tau \geq 0$, $\xi \in [0, 1]$ be such a homotopy that λ_{ξ} are geodesic rays (not necessarily timelike) emitted from a single point $p \equiv \lambda_{\xi}(0) \notin \tilde{H}'$ and lying in $\tilde{H} - \tilde{\Theta}$. Denote by $h_{\wedge}(\xi)$ and $h_{\vee}(\xi)$ the intersections of λ_{ξ} with $\mathcal{H}'_{\wedge}$ and $\mathcal{H}'_{\vee}$, respectively. The corresponding values of τ we shall denote by $\tau_{\wedge}$ and $\tau_{\vee}$:

$$\tau_{\wedge, \vee}(\xi) \equiv \lambda_{\xi}^{-1}[h_{\wedge, \vee}(\xi)].$$

Since both $\tilde{H}$ and $\tilde{H}'$ are convex the segment of λ_{ξ} between $h_{\wedge}(\xi)$ and $h_{\vee}(\xi)$ (when they both exist) lies in $\tilde{H}'$. Moreover, as λ_{ξ} does not intersect $\tilde{\Theta}$ it follows from (19) that the segment contains a point of $\tilde{H}'$ and hence (see proposition 8) the whole segment $\lambda_{\xi}[(\tau_{\wedge}, \tau_{\vee})]$ lies in $\tilde{H}'$. Which means, in particular, that λ_{ξ} do not intersect $\mathcal{H}'_{\wedge}$ and $\mathcal{H}'_{\vee}$ except in $h_{\wedge, \vee}(\xi)$.

36. Proposition. If λ_{ξ_0} with $\xi_0 \in [0, 1]$ intersects both $\mathcal{H}'_{\wedge}$ and $\mathcal{H}'_{\vee}$, then so do all the other λ_{ξ} and the functions $\tau_{\vee, \wedge}(\xi)$ are continuous.

Proof. Consider the (non-empty by assumption) set $\Xi \subset [0, 1]$ defined by

$$\Xi \equiv \{\xi: \exists h_{\vee}(\xi), h_{\wedge}(\xi)\}.$$

Suppose $\xi_1 \in \Xi$. Then

$$\begin{aligned} \lambda_{\xi_1}(\tau) &\in \tilde{H}_{\wedge}, & \tau &\in [\tau_{\wedge} - \epsilon, \tau_{\wedge} + \epsilon] \\ \lambda_{\xi_1}(\tau_{\wedge} - \epsilon) &\in \tilde{H}_{\wedge} - \overline{\tilde{H}'_{\wedge}}, & \lambda_{\xi_1}(\tau_{\wedge} + \epsilon) &\in \tilde{H}'_{\wedge} \end{aligned}$$

assuming that ϵ is sufficiently small and that (for definiteness) $\tau_{\wedge} < \tau_{\vee}$. Obviously this holds also for any ξ sufficiently close to ξ_1 . Consequently all λ_{ξ} with such ξ also intersect $\text{Bd } \tilde{H}'_{\wedge}$ and hence (recall that λ_{ξ} do not pass through $\tilde{\Theta}$) $\mathcal{H}'_{\wedge}$. The same is true for $\mathcal{H}'_{\vee}$. So Ξ is open.

Now let ξ_2 be a limit point of Ξ . $\mathcal{H}'_{\wedge}$ is compact, so it contains the limit point h of $h_{\wedge}(\xi)$:

$$\mathcal{H}'_{\wedge} \ni h = \lim_{\xi \rightarrow \xi_2} h_{\wedge}(\xi) = \lim_{\xi \rightarrow \xi_2} \lambda_{\xi}(\tau_{\wedge}) = \lambda_{\xi_2}(\tau_*). \quad (20)$$

Similarly λ_{ξ_2} must intersect $\mathcal{H}'_{\vee}$. So, Ξ is closed and hence is equal to $[0, 1]$. From (20) it follows also that $\tau_{\wedge, \vee}(\xi_2) = \tau_* = \lim_{\xi \rightarrow \xi_2} \tau_{\wedge, \vee}(\xi)$ whence $\tau_{\wedge, \vee}(\xi)$ are continuous. $\square$

37. Remark. As was mentioned in remark 26, N is an area in a covering of $\tilde{H} - \tilde{\Theta}$. Because this area is bounded, not *all* curve in $\tilde{H} - \tilde{\Theta}$ can be lifted to a continuous curve

in N . It is easy to check, however, that if a curve $\tilde{\gamma}: [0, 1] \rightarrow \tilde{H}$ satisfies the following condition

$$\tilde{\gamma}(0) \notin \overline{\tilde{H}'}, \quad \tilde{\gamma} \cap \mathcal{H}'_\Lambda = \emptyset, \text{ or } \tilde{\gamma} \cap \mathcal{H}'_\nu = \emptyset, \quad (21)$$

then there is a unique curve $\gamma \subset N$ connecting $\pi^{-1}(\tilde{\gamma}(0))$ with $\pi^{-1}(\tilde{\gamma}(1))$ and satisfying $\pi(\gamma) = \tilde{\gamma}$. It is this γ that we understand by $\pi^{-1}(\tilde{\gamma})$ from now on.

38. Proposition. Let $\lambda_\xi(\tau)$ be the homotopy from proposition 36, $\tilde{\mu}(\xi) \equiv \lambda_\xi(1)$, and $\tilde{\gamma}_\xi(\tau) \equiv \lambda_\xi(\tau)|_{\tau \in [0,1]}$. Suppose that $\tilde{\mu}$ does not intersect $\tilde{\mathcal{S}}_-$ and that $\tilde{\mu}(0) \notin \tilde{H}'$. If $\tilde{\gamma}_0$ satisfies (21) then so does $\tilde{\gamma}_1$.

Proof. Suppose for contradiction that $\tilde{\gamma}_1$ (and hence λ_1) intersects both $\mathcal{H}'_\Lambda$ and $\mathcal{H}'_\nu$:

$$\tau_\Lambda(1), \tau_\nu(1) \leq 1.$$

Then (by proposition 36) so does λ_0 *in contrast to* $\tilde{\gamma}_0$, which is only possible if $\lambda_0(1)$ lies on λ_0 closer to p than $h_\Lambda(0)$ (and thus also than $h_\nu(0)$ assuming as before that $\tau_\Lambda < \tau_\nu$). So,

$$\tau_\Lambda(0), \tau_\nu(0) > 1$$

and there must exist ξ' and ξ'' :

$$\xi' \equiv \max\{\xi \mid \tau_\Lambda(\xi) \geq 1\}, \quad \xi'' \equiv \min\{\xi \mid \xi > \xi', \tau_\nu(\xi) \leq 1\}, \quad 0 < \xi' < \xi'' < 1.$$

The (open) segment of $\tilde{\mu}$ between ξ' and ξ'' is a continuous curve lying within $\tilde{H}'$ (since $\tau_\Lambda < 1 < \tau_\nu$ here) and connecting $\tilde{H}'_\Lambda$ with $\tilde{H}'_\nu$ without intersecting $\tilde{\mathcal{S}}_-$, which is impossible. □

5. Proof of the theorem

For any region U denote by $\mathcal{V}(U)$ the *causality violating subset* of U , that is the set of all points p satisfying $J_U^+(p) \cap J_U^-(p) \neq \{p\}$. Note that $\mathcal{V}(U)$ is determined by the causal structure of U , but not by that of the ambient spacetime $U' \supset U$ (if it exists), in the sense that generally $\mathcal{V}(U) \subsetneq \mathcal{V}(U') \cap U$.

39. Proposition. Any extendible LCC spacetime M has an LCC extension whose causality violating subset is $\mathcal{V}(M)$.

Proof. We shall prove that an extension $M_\mathcal{N}$, which (as was shown in section 3) can be built for any extendible LCC M , satisfies the requirement of the proposition, i. e. that $M_\mathcal{N}$ is LCC and $\mathcal{V}(M_\mathcal{N}) = \mathcal{V}(M)$. The latter follows immediately from remark 25 (since $M_\mathcal{N}$ can be isometrically immersed into M_Δ) and to prove the former suppose that the assertion is false. Then there exists a future-directed curve $\lambda \subset D$ from a to b such that

$$a, b \in D_M, \quad \lambda \not\subset D_M.$$

Without loss of generality we may assume that λ does not intersect $\mathcal{S}_\infty$ (and thus has no roots). By proposition 33 this implies that the ends of λ lie either both in M , or both in $H'_{\mathcal{N}\wedge}$, or both in $K_\diamond$. But M and $H'_{\mathcal{N}}$ are LCC, which implies that the ends of λ cannot lie in a connected component of $D \cap M$, or $D \cap H'_{\mathcal{N}}$ and thus by proposition 34 in $D_M \cap M$, or $D_M \cap H'_{\mathcal{N}}$. So, $a, b \in K_\diamond$.

Choose a and b so that $\pi(a), \pi(b) \notin \overline{\tilde{H}'}$ (it always can be done). We want to show that they can be connected by a *timelike* curve — it will be the geodesic $\tilde{\gamma}_1$ — lying in $D \cap K_\diamond$. To this end let us, first, pick a curve $\mu(\xi): [0, 1] \rightarrow D \cap K_\diamond$ connecting a and b (its existence is guaranteed by proposition 34). μ can always be chosen (cf. proposition 33) so that for some $x, y \in D$

$$\mu \subset \langle x, y \rangle_D.$$

Consider the geodesics $\tilde{\gamma}_\xi \subset \tilde{H}$ with $\tilde{\gamma}_\xi(0) = \tilde{\mu}(0)$ and $\tilde{\gamma}_\xi(1) = \tilde{\mu}(\xi)$ (as usual $\tilde{\mu} \equiv \pi(\mu)$). Let Ξ be the set of all ξ such that

$$\tilde{\gamma}_{\xi'} \text{ satisfies (21), } \gamma_{\xi'} \equiv \pi^{-1}(\tilde{\gamma}_{\xi'}) \subset \langle x, y \rangle_D \quad \forall \xi' \leq \xi. \quad (22)$$

Ξ is evidently open and (as small ξ obviously lie in it) non-empty. To see that it is also closed consider its limit point ξ_c

$$\xi_n \rightarrow \xi_c, \quad \xi_n \in \Xi.$$

By construction none of $\tilde{\gamma}_{\xi'}$ with $\xi' < \xi_c$ meets $\tilde{\Theta}$. It is easy to check that neither does $\tilde{\gamma}_{\xi_c}$. Whence by proposition 38 the condition (21) holds for ξ_c . Consequently there exists a geodesic $\gamma_{\xi_c} \equiv \pi^{-1}(\tilde{\gamma}_{\xi_c})$, which (being the limit of γ_{ξ_n}) lies in $\langle x, y \rangle_D \subset D$ and moreover (since a and b belong to a convex subset $\langle x, y \rangle_D$ of D) in $\langle x, y \rangle_D$. Thus the whole condition (22) holds for ξ_c and Ξ is closed. So, $\Xi = [0, 1]$.

We have proved, thus, that $\gamma_1 \subset D \cap N$. Besides, γ_1 is timelike (since b is connected to a with the timelike $\lambda \subset D$) and $n_\pm[\gamma_1] = 0$ (since $n_\pm[\lambda] = 0$), whence $\gamma_1 \subset K_\diamond$. So, in $D \cap K_\diamond$ a timelike curve from a to b exists indeed. Which yields

$$\lambda(\epsilon), \lambda(1 - \epsilon) \in D_K, \quad (23a)$$

$$\lambda([\epsilon, 1 - \epsilon]) \subset D_1, \quad \lambda([\epsilon, 1 - \epsilon]) \not\subset D_K, \quad (23b)$$

$$D_1 \cap \mathcal{S}_\infty = \emptyset, \quad (23c)$$

where $D_1 \equiv \langle a, b \rangle_D$, D_K is a component of $D_1 \cap K_\diamond$, and ϵ is chosen appropriately small. To see that (23) is, in fact, impossible consider the region $D_1 \cup K_\diamond$ of the spacetime M^e as a separate spacetime and extend it to a spacetime H^e by gluing $\tilde{H}$ to it [(23c) ensures that the resulting manifold is Hausdorff):

$$H^e \equiv (D_1 \cup K_\diamond) \cup_{\varpi_K} \tilde{H}$$

(in other words H^e is what results when the ‘hole’ in $K_\diamond$, which appeared when $D_1 \cup K_\diamond$ was cut out of M^e , is ‘closed up’ by gluing its edges (the former $\mathcal{S}_-$ and $\mathcal{S}_+$) together). H^e is an extension of $\tilde{H}$ and D_1 is (see remark 16) its diamond subset. So (23a,b) contradict the local causal convexity of $\tilde{H}$.

□

40. Proposition. Any extendible LCC spacetime M has a maximal extension $M^{\max}$ with $\mathcal{V}(M^{\max}) = \mathcal{V}(M)$.

Proof. Consider the set $\mathbf{E}$ of all pairs (V, ζ) , where ζ is an isometric imbedding of M into V (we normally shall not distinguish $\zeta(M)$ and M) and V is an LCC extension of M with $\mathcal{V}(V) = \mathcal{V}(M)$. Pick a point $p \in M$ and introduce the following order relation in $\mathbf{E}$ (cf. [7]). We write $(V_1, \zeta_1) \leq (V_2, \zeta_2)$ if there exists an isometric imbedding $\vartheta_{12}: V_1 \rightarrow V_2$ satisfying the conditions:

$$\zeta_2^{-1} \circ \vartheta_{12} \circ \zeta_1(p) = p, \quad d(\zeta_2^{-1} \circ \vartheta_{12} \circ \zeta_1)|_p = \text{id}.$$

It is easy to check that in agreement with its notation $\leq$ is a partial order (which it would not be if we relax any of the two conditions). Let $\{(V_\alpha, \zeta_\alpha)\}$ be a chain with respect to $\leq$ and let

$$V = \left(\bigcup_{\alpha} V_{\alpha} \right) \setminus \sim,$$

where the equivalence is defined as follows:

$$x \sim y \Leftrightarrow \exists \alpha_1, \alpha_2: \vartheta_{\alpha_1 \alpha_2}(x) = y, \text{ or } \vartheta_{\alpha_1 \alpha_2}(y) = x.$$

Let us further introduce the following notations: V^e is an extension of V ; D is a diamond subset of V^e ; D_V is a connected component of $V^e \cap D$; $\varphi \subset D_V$ is a curve between points a and b ; $\lambda_{ab} \subset D$ is a timelike curve from a to b ; and $\ell \subset V$ is a closed causal curve.

Since the curves φ and ℓ are compact we can choose a finite number of V_α : $\{V_{\alpha_n}\}$, $n = 1, \dots, N$ so that (from now on we do not distinguish V_α and their images in V)

$$\varphi, \ell \subset \bigcup_n V_{\alpha_n} \subset V_{\alpha_0}. \quad (24)$$

Along with the whole φ the points a, b lie in a connected component of $V_{\alpha_0} \cap D$ and (since V_{α_0} is LCC) so does λ_{ab} . Hence V is LCC. Also $\mathcal{V}(V)$ is equal to $\mathcal{V}(M)$, since otherwise [by (24)] $\mathcal{V}(V_{\alpha_0})$ would not be. Finally, C holds in V since it is covered by the open sets V_{α_n} where it does. So, $(V, \zeta_1) \in \mathbf{E}$ and obviously $(V, \zeta_1) = \sup\{(V_\alpha, \zeta_\alpha)\}$. Which means that $\leq$ is an inductive order in $\mathbf{E}$. It follows then by the Zorn lemma that M has an LCC extension $M^{\max}$ with $\mathcal{V}(M^{\max}) = \mathcal{V}(M)$ which cannot be extended to any larger spacetime with such properties. By proposition 39 $M^{\max}$ is maximal. □

Proof of the theorem. Consider the set of all possible extensions of U of the form $I_{U'}^-(U)$, where U' is an arbitrary spacetime. By repeating the procedure employed in proposition 40 it can be shown that U can be imbedded into a maximal element — let us call it M — of that set. $M = I_{M^e}^-(U)$ in any extension M^e of M . So M is obviously LCC (since a past-directed curve even cannot leave M) and (tautologically) $\mathcal{V}(M) \subset I_{M^e}^-(U)$. Which being combined with proposition 40 proves the theorem. □

Acknowledgments

I wish to thank R. R. Zapatrin for useful discussion and Starlab NV for wonderful conditions which I enjoyed while working on this paper.

References

- [1] D. H. Politzer 1992 *Phys. Rev. D* **46** 4470
- [2] M. S. Morris, K. S. Thorne, and U. Yurtsever 1988 *Phys. Rev. Lett.* **61** 1446
- [3] M. Visser 1996 *Lorentzian Wormholes* (NY: AIP Press)
P. J. Nahin 1999 *Time Machines* (NY: Springer-Verlag)
J. Earman. and C. Smeenk *Reverberations of the Shaky Game* (eds R. Jones and P. Ehrlich, to be published by University of Chicago Press)
- [4] S. W. Hawking 1992 *Phys. Rev. D* **46** 603
S. Krasnikov 1998 *Class. Quantum Grav.* **15** 997
- [5] P. T. Chruściel and A. D. Rendall 1995 *Ann. Phys., NY* **242** 349
- [6] S. Krasnikov 2002 *Phys. Rev. D* **65** 064013
- [7] S. W. Hawking and G. F. R. Ellis 1973 *The Large Scale Structure of Spacetime* (Cambridge: Cambridge University Press, England)
- [8] B. O'Neill *Semi-Riemannian Geometry* (Academic Press, 1983)
- [9] R. Penrose *Techniques of Differential Topology in Relativity*, (SIAM, 1983).
- [10] U. Yurtsever 1994 *Class. Quantum Grav.* **11** 1013
- [11] S. V. Krasnikov 1999 *Phys. Rev. D* **59** 024010